

Pretzelosity distribution function $h_{1T}^\perp$

H. Avakian,¹ A. V. Efremov,² P. Schweitzer,^{3,4} F. Yuan^{5,6}

¹ *Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, U.S.A.*

² *Joint Institute for Nuclear Research, Dubna, 141980 Russia*

³ *Inst. f. Theoret. Physik II, Ruhr-Universität Bochum, D-44780 Bochum, Germany*

⁴ *Departement of Physics, University of Connecticut, Storrs, CT 06269, U.S.A.*

⁵ *RIKEN BNL Research Center, Building 510A, BNL, Upton, NY 11973, U.S.A.*

⁶ *Nuclear Science Division, Berkeley National Laboratory, Berkeley, CA 94720, U.S.A.*

The 'pretzelosity' distribution $h_{1T}^\perp$ is discussed. Theoretical properties, model results, and perspectives to access experimental information on this leading twist, transverse momentum dependent parton distribution function are reviewed. Its relation to helicity and transversity distributions is highlighted.

Keywords: semi-inclusive deep inelastic scattering (SIDIS), single spin asymmetry (SSA), transverse momentum dependent distribution function (TMD)

1. Introduction

SIDIS allows to access information on TMDs that are defined in terms of light-front correlators¹⁻⁶ (with process-dependent paths⁷⁻¹⁰)

$$\phi(x, \vec{p}_T)_{ij} = \int \frac{dz^- d^2 \vec{z}_T}{(2\pi)^3} e^{ipz} \langle N | \bar{\psi}_j(0) \mathcal{W}(0, z, \text{path}) \psi_i(z) | N \rangle \Big|_{\substack{z^+ = 0 \\ p^+ = xP^+}} \quad (1)$$

where light-cone components $p^\pm = (p^0 \pm p^3)/\sqrt{2}$ are along the virtual photon momentum, and transverse vectors like $\vec{p}_T$ are perpendicular to it. Different TMDs are given by traces of the correlator (1) with specific γ -matrices. There are 8 leading-twist TMDs, i.e. they give rise to effects that are not power suppressed in the hard scale, and each of them contains independent information about the nucleon structure. All leading-twist TMDs can be accessed unambiguously in SIDIS with polarized leptons and nucleons by observing the azimuthal distributions of produced hadrons.

In this note we will focus on the leading-twist TMD pretzelosity $h_{1T}^{\perp a}$, on which interesting results have been obtained from model calculations. This TMD is responsible for a SSA $\propto \sin(3\phi - \phi_S)$ in SIDIS that could be accessed in experiments — most promisingly at Jefferson Lab.

2. Properties of $h_{1T}^{\perp a}$

Let us list briefly, what we know about the pretzelosity distribution.

- I. It can be projected out from the correlator (1) by tracing it with $i\sigma^{j+}\gamma_5$ where it appears as the coefficient of the structure $S_T^k(p_T^j p_T^k - \frac{1}{2} \vec{p}_T^2 \delta^{jk})$, and it has a probabilistic interpretation.⁶
- II. It requires nucleon wave-function components with two units orbital momentum difference,¹¹ and 'measures' the deviation of the 'nucleon shape' from a sphere.¹² (That is why it is called 'pretzelosity'!)
- III. It is expected to be suppressed at small and large^{11,13,14} x with respect to parton distributions like $f_1^a(x)$, $g_1^a(x)$, $h_1^a(x)$.
- IV. It satisfies the positivity condition¹⁵ $2|h_{1T}^{\perp(1)a}(x)| \leq f_1^a(x) - g_1^a(x)$.
The above and Soffer bound¹⁶ imply: $|h_{1T}^{\perp(1)a}(x)| + |h_1^a(x)| \leq f_1^a(x)$.
- V. In the limit of a large number of colors N_c in QCD it was shown that¹⁷ $(h_1^{\perp u} + h_1^{\perp d})/(h_1^{\perp u} - h_1^{\perp d}) \sim \mathcal{O}(1/N_c)$ for $xN_c \sim \mathcal{O}(1)$ and $p_T \sim \mathcal{O}(1)$. The same pattern is predicted also for antiquarks.¹⁷
- VI. It was observed in the bag model¹⁸ that $h_{1T}^{\perp(1)q}(x) = g_1^q(x) - h_1^q(x)$ which is confirmed in many¹⁸⁻²² (but not all^{22,23}) models.
- VII. In simple (spectator-type) models, it has been related to chirally odd generalized parton distributions.²²
- VIII. In SIDIS with unpolarized electrons (U) and transversely polarized nucleons (T) it gives rise (in combination with Collins fragmentation function^{24,25}) to an azimuthal modulation of the produced hadrons proportional to $\sin(3\phi - \phi_S)$. Here ϕ (ϕ_S) is the azimuthal angle of the produced hadron (nucleon polarization vector) with respect to the virtual photon. The corresponding SSA is given by

$$A_{UT}^{\sin(3\phi - \phi_S)} = \frac{C_{\text{Gauss}} \sum_a e_a^2 x h_{1T}^{\perp(1)a}(x) H_1^{\perp(1/2)a}(z)}{\sum_a e_a^2 x f_1^a(x) D_1^a(z)} \quad (2)$$

with $h_{1T}^{\perp(1)}(x) = \int d^2 \vec{p}_T \frac{\vec{p}_T^2}{2M_N^2} h_{1T}^{\perp}(x, \vec{p}_T^2)$ the 'transverse moment' of pretzelosity. Unless the DIS counts are weighted with adequate powers of transverse hadron momenta⁵ it is necessary to assume a model for the distribution of transverse parton momenta. In (2) the Gauss model is assumed. The factor C_{Gauss} contains the dependence on Gauss model parameters and is a slowly varying function of these parameters that can be well approximated for practical purposes¹⁸ by its maximum $C_{\text{Gauss}} \leq C_{\text{max}} = 3/(2\sqrt{2})$. Extractions of the (1/2)-moment of the Collins function $H_1^{\perp(1/2)}(z) = \int d^2 \vec{K}_T \frac{|\vec{K}_T|}{2z m_h} H_1^{\perp}(z, \vec{K}_T^2)$ from data²⁶⁻³⁰ were reported elsewhere.³¹⁻³³

3. Pretzelosity in the bag model

In this Section we review the pretzelosity calculation¹⁸ in the MIT bag model, in which the nucleon consists of 3 non-interacting quarks confined in a spherical cavity.³⁴⁻³⁷ The momentum space wave function is given by

$$\varphi_m(\vec{k}) = i\sqrt{4\pi}NR_0^3 \begin{pmatrix} t_0(|\vec{k}|)\chi_m \\ \vec{\sigma} \cdot \hat{k} t_1(|\vec{k}|)\chi_m \end{pmatrix}, \quad (3)$$

with $N = \omega^{3/2}(2R_0^3(\omega - 1)\sin^2\omega)^{-1/2}$, $\hat{k} = \vec{k}/|\vec{k}|$. We fix the bag radius R_0 in terms of the proton mass M_N as $R_0M_N = 4\omega$ with $\omega \approx 2.04$ the lowest root of the bag eigen-equation. Finally $t_i(\kappa) = \int_0^1 u^2 du j_i(uR_0\kappa)j_i(u\omega)$ where j_i are spherical Bessel functions. The bag model wave function (3) contains both S (represented by t_0) and P (represented by t_1) waves.

With the above wave functions, one obtains the following results¹⁸

$$f_1(x, k_\perp) = A \left[t_0^2 + 2t_0t_1 \frac{k_z}{k} + t_1^2 \right] \quad (4)$$

$$g_1(x, k_\perp) = A \left[t_0^2 + 2t_0t_1 \frac{k_z}{k} + t_1^2 \left(\frac{2k_z^2}{k^2} - 1 \right) \right] \quad (5)$$

$$h_1(x, k_\perp) = A \left[t_0^2 + 2t_0t_1 \frac{k_z}{k} + t_1^2 \frac{k_z^2}{k^2} \right] \quad (6)$$

$$h_{1T}^\perp(x, k_\perp) = A \left[-2 \frac{M_N^2}{k^2} t_1^2 \right] \quad (7)$$

with $A = 16\omega^4/[\pi^2(\omega - 1)j_0^2(\omega)M_N^2]$ and the flavor dependence given by (we assume an $SU(6)$ spin-flavor symmetric proton wave function):

$$\begin{aligned} f_1^q(x, k_\perp) &= N_q f_1(x, k_\perp), \quad N_u = 2, \quad N_d = 1 \\ g_1^q(x, k_\perp) &= P_q g_1(x, k_\perp), \quad P_u = \frac{4}{3}, \quad P_d = -\frac{1}{3}. \end{aligned} \quad (8)$$

The flavor dependence of h_1^q and $h_{1T}^{\perp q}$ is analog to g_1^q . The momenta k_z and k are defined as $k_z = xM_N - \omega/R_0$, and $k = \sqrt{k_z^2 + k_\perp^2}$.

Since $h_{1T}^\perp \propto t_1^2$ it is proportional to the square of the P -wave¹¹ and thus sensitive to the quark orbital angular momentum in the proton. The results (4-8) satisfy¹⁸ the positivity conditions in point III, and are consistent¹⁸ with the predictions from the large- N_c limit discussed in point IV of Sec. 2.

From the results in Eqs. (4-7) we find that out of the 4 TMDs $f_1(x, k_\perp)$, $g_1(x, k_\perp)$, $h_1(x, k_\perp)$, $h_{1T}^{\perp(1)}(x, k_\perp) \equiv k_\perp^2/(2M_N^2) h_{1T}^\perp(x, k_\perp)$ only 2 are linearly independent. In QCD the different TMDs are, of course, all independent of each other. But in the bag model all TMDs are expressed in terms of t_0 and t_1 in Eq. (3), which naturally gives rise to *bag model relations*




Fig. 1. The parton distribution function $h_{1T}^{\perp q}(x)$ vs. x from the bag model¹⁸ in comparison to $f_1^q(x)$ and $h_1^q(x)$ from this model. The results refer to a low scale.³⁶ Notice that $h_{1T}^{\perp q}(x)$ is not constrained by positivity.

among different TMDs. The most interesting (and possibly more general, see Sec. 4) relation is¹⁸

$$h_{1T}^{\perp(1)q}(x, k_{\perp}) = g_1^q(x, k_{\perp}) - h_1^q(x, k_{\perp}). \quad (9)$$

Figs. 1a and 1b show results for $h_{1T}^{\perp q}(x) = \int d^2\vec{k}_{\perp} h_{1T}^{\perp q}(x, k_{\perp})$ at the low bag model scale. The pretzelosity distributions $h_{1T}^{\perp q}(x)$ have opposite signs compared to transversity and are larger than $h_1^q(x)$ in the bag model, even larger than $f_1^q(x)$. However, $h_{1T}^{\perp q}(x)$ is not constrained by positivity bounds.

The bag¹⁸ (and spectator¹⁹) model results satisfy the positivity bounds for $h_{1T}^{\perp(1)q}(x)$ in point III of Sec. 2. In Fig. 2 the results for $h_{1T}^{\perp(1)q}(x)$ from both models are compared. We observe good qualitative agreement.

The moduli of the transverse moments $h_{1T}^{\perp(1)q}(x)$ of the pretzelosity distribution functions are about 3 or more times smaller than those of the transversity distribution functions $h_1^q(x)$ with the exception of very small- x where both models are strictly speaking not applicable.¹⁸




Fig. 2. The $h_{1T}^{\perp(1)q}(x)$ vs. x at low scale from bag model¹⁸ and spectator model.¹⁹

4. How general is the relation in Eq. (9) ?

Eq. (9) is remarkable from the point of view of the observation that³⁸ 'the difference of $g_1^q(x)$ and $h_1^q(x)$ is a measure for relativistic effects in nucleon'. This difference is just the transverse moment of pretzelosity!

It is clear that this relation cannot be strictly valid in QCD, where all TMDs are independent. However, could it nevertheless allow to gain a reasonable estimate for $h_{1T}^{\perp(1)}(x)$ in terms of transversity and helicity? Until clarified by experiment, we can address this question only in models.

Interestingly, the relation (9) does not hold only in the bag model,¹⁸ but is found¹⁸ to be satisfied also in the spectator model.¹⁹ In fact, it was conjectured¹⁸ that (9) could hold in a large class of *relativistic quark models*, which was subsequently confirmed in the constituent quark model²⁰ and the relativistic model of the proton.²¹ But (9) is not satisfied in a different than¹⁹ version of the spectator model.²³

The limitations of (9) are nicely illustrated in the 'quark target model' where in addition to quarks there are also gluons, and (9) is not satisfied.²² Thus, the explicit inclusion of gluon degrees of freedom spoils this relation. Of course, as stressed above, we do not expect (9) to be valid in QCD.

It would be interesting to know the necessary and sufficient conditions for the relation (9) to hold in a model.

5. Preliminary COMPASS data & prospects at JLab

At COMPASS the $\sin(3\phi - \phi_S)$ and other SSAs were measured on a deuteron target.³⁹ By saturating the positivity bound for $h_{1T}^{\perp(1)}(x)$ (point III in Sec. 2) we estimated¹⁸ the maximum effect for $A_{UT}^{\sin(3\phi - \phi_S)}$. For that information




Fig. 3. The transverse target SSA $A_{UT, \pi}^{\sin(3\phi - \phi_S)}$ for deuteron estimated on the basis of the positivity bound vs. preliminary COMPASS data.³⁹

on Collins effect^{32,33} and parameterizations^{40,41} were used. The results are shown in Fig. 3 and compared to the preliminary data.³⁹

At small x the preliminary data favor that pretzelosity does not reach the bound. Whether due to the expected suppression at small x or opposite signs of u and d -flavors, see Sec. 2, cannot be concluded.

The important observation is that preliminary COMPASS data³⁹ do not exclude a sizeable effect in the region $x > 0.1$, see Fig. 3, where JLab can measure with precision. This is demonstrated in Fig. 4 showing the π^+ SSA from a proton target in the kinematics of CLAS with 12 GeV beam upgrade (with error projections for 2000 hours run time⁴²).

It could be even more promising to look at SSAs due to Collins effect, like $A_{UT}^{\sin(3\phi-\phi_S)}$, in kaon production. The statistics for kaon production is lower than for pion production, but the SSA might be larger as it is suggested by a model⁴³ of the Collins function. With a RICH detector at CLAS the kaon SSAs could be measured in the valence- x region.⁴⁴




Fig. 4. SSA $A_{UT}^{\sin(3\phi-\phi_S)}$ in π^+ production from proton in the kinematics of CLAS 12 as function of x (error projections from⁴²). The shaded areas indicate the range allowed by positivity. The bag¹⁸ and spectator¹⁹ models predict a negative SSA.

6. Conclusions

We reviewed the bag model calculation of pretzelosity.¹⁸ An interesting observation is the relation (9) which connects helicity and transversity distributions to $h_{1T}^{\perp(1)}(x)$, and is valid in many¹⁸⁻²² (though not all^{22,23}) models, but not in QCD where all TMDs are independent.⁴⁵ Nevertheless (9) could turn out a useful approximation. In view of the numerous novel TMDs, well-motivated approximations are welcome.⁴⁶

In the bag model $h_{1T}^{\perp u}$ is proportional to *minus* the square of the p -wave component of the nucleon wave function, and therefore manifestly negative ($h_{1T}^{\perp d}$ has opposite sign dictated by SU(6) symmetry, and large N_c ¹⁷). As a consequence of (9) we have $|h_1^q(x)| > |g_1^q(x)|$. This is found in models.^{47,48}

Forthcoming analyzes and experiments at COMPASS, HERMES and JLab^{42,49,50} will provide valuable information on the pretzelosity distribution function, and deepen our understanding of the nucleon structure.

Acknowledgements. We thank the organizers of the *Transversity 2008* for the opportunity to present our work supported by: BMBF, German-Russian collaboration (DFG-RFFI, 436 RUS 113/881/0), the EIIIHP project RII3-CT-2004-506078, the Grants RFBR 06-02-16215 and 07-02-91557, RF MSE RNP.2.2.2.2.6546 (MIREA), the Heisenberg-Landau Program of JINR, the contract DE-AC05-06OR23177, RIKEN, Brookhaven National Laboratory, and the U.S. Department of Energy DE-AC02-98CH10886.

References

1. J. C. Collins and D. E. Soper, Nucl. Phys. B **193**, 381 (1981) [Erratum-ibid. B **213**, 545 (1983)].
2. J. C. Collins, D. E. Soper and G. Sterman, Nucl. Phys. B **250**, 199 (1985).
3. X. D. Ji, J. P. Ma and F. Yuan, Phys. Rev. D **71**, 034005 (2005).
X. D. M. Ji, J. P. M. Ma and F. Yuan, Phys. Lett. B **597**, 299 (2004).
4. J. C. Collins and A. Metz, Phys. Rev. Lett. **93**, 252001 (2004).
5. D. Boer and P. J. Mulders, Phys. Rev. D **57**, 5780 (1998).
6. P. J. Mulders and R. D. Tangerman, Nucl. Phys. B **461** (1996) 197 [Erratum-ibid. B **484** (1997) 538].
7. S. J. Brodsky, D. S. Hwang and I. Schmidt, Phys. Lett. B **530**, 99 (2002); Nucl. Phys. B **642**, 344 (2002).
8. J. C. Collins, Phys. Lett. B **536**, 43 (2002) [arXiv:hep-ph/0204004].
9. A. V. Belitsky, X. Ji and F. Yuan, Nucl. Phys. B **656**, 165 (2003).
X. D. Ji and F. Yuan, Phys. Lett. B **543**, 66 (2002).
D. Boer, P. J. Mulders and F. Pijlman, Nucl. Phys. B **667**, 201 (2003).
10. I. O. Cherednikov and N. G. Stefanis, Phys. Rev. D **77**, 094001 (2008);
I. O. Cherednikov and N. G. Stefanis, Nucl. Phys. B **802**, 146 (2008).
11. M. Burkardt, arXiv:0709.2966 [hep-ph].
12. G. A. Miller, Phys. Rev. C **76**, 065209 (2007).
13. H. Avakian, S. J. Brodsky, A. Deur and F. Yuan, Phys. Rev. Lett. **99**, 082001 (2007).
14. S. J. Brodsky and F. Yuan, Phys. Rev. D **74** (2006) 094018.
15. A. Bacchetta, M. Boglione, A. Henneman and P. J. Mulders, Phys. Rev. Lett. **85**, 712 (2000).
16. J. Soffer, Phys. Rev. Lett. **74** (1995) 1292.
17. P. V. Pobylitsa, arXiv:hep-ph/0301236.
18. H. Avakian, A. V. Efremov, P. Schweitzer and F. Yuan, arXiv:0805.3355.
19. R. Jakob, P. J. Mulders and J. Rodrigues, Nucl. Phys. A **626**, 937 (1997).
20. B. Pasquini, S. Cazzaniga and S. Boffi, arXiv:0806.2298 [hep-ph].
B. Pasquini, arXiv:0807.2825 [hep-ph].
21. A. V. Efremov, P. Schweitzer, O. V. Teryaev and P. Zavada, in preparation (P. Zavada, talk at *SPIN Praha 2008*, Prague, 20-26 July 2008).
22. S. Meissner, A. Metz and K. Goeke, Phys. Rev. D **76**, 034002 (2007).
23. A. Bacchetta, F. Conti and M. Radici, arXiv:0807.0323 [hep-ph].
24. J. C. Collins, Nucl. Phys. B **396**, 161 (1993).

25. A. V. Efremov, L. Mankiewicz and N. A. Tornqvist, Phys. Lett. B **284** (1992) 394. J. C. Collins, S. F. Heppelmann and G. A. Ladinsky, Nucl. Phys. B **420** (1994) 565.
26. A. Airapetian *et al.* [HERMES Coll.], Phys. Rev. Lett. **94**, 012002 (2005).
27. V. Y. Alexakhin *et al.* [COMPASS Coll.], Phys. Rev. Lett. **94**, 202002 (2005).
28. M. Diefenthaler [HERMES Coll.], AIP Conf. Proc. **792** (2005) 933, arXiv:0706.2242 and arXiv:hep-ex/0612010.
29. E. S. Ageev *et al.* [COMPASS Coll.], Nucl. Phys. B **765** (2007) 31
30. M. Alekseev *et al.* [COMPASS Coll.], arXiv:0802.2160 [hep-ex].
31. W. Vogelsang and F. Yuan, Phys. Rev. D **72** (2005) 054028.
32. A. V. Efremov, K. Goeke and P. Schweitzer, Phys. Rev. D **73**, 094025 (2006).
33. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin and C. Turk, Phys. Rev. D **75**, 054032 (2007).
34. A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. D **9**, 3471 (1974). R. L. Jaffe, Phys. Rev. D **11**, 1953 (1975).
35. L. S. Celenza and C. M. Shakin, Phys. Rev. C **27**, 1561 (1983) [Erratum-ibid. C **39**, 2477 (1989)].
36. M. Stratmann, Z. Phys. C **60**, 763 (1993).
37. F. Yuan, Phys. Lett. B **575**, 45 (2003).
38. R. L. Jaffe and X. D. Ji, Nucl. Phys. B **375** (1992) 527, and Phys. Rev. Lett. **67**, 552 (1991).
39. A. Kotzinian [on behalf of the COMPASS Coll.], arXiv:0705.2402 [hep-ex].
40. M. Glück, E. Reya and A. Vogt, Eur. Phys. J. C **5**, 461 (1998). M. Glück, E. Reya, M. Stratmann and W. Vogelsang, Phys. Rev. D **63**, 094005 (2001).
41. S. Kretzer, Phys. Rev. D **62** (2000) 054001.
42. H. Avakian, *et al.*, JLab LOI 12-06-108 (2008).
43. A. Bacchetta, L. P. Gamberg, G. R. Goldstein and A. Mukherjee, Phys. Lett. B **659**, 234 (2008) [arXiv:0707.3372 [hep-ph]].
44. CLAS 12 Rich Detector Workshop, 28-29 January 2008, Jefferson Lab, Newport News, VA, U.S.A. <http://conferences.jlab.org/CLAS12>
45. K. Goeke, A. Metz and M. Schlegel, Phys. Lett. B **618**, 90 (2005).
46. H. Avakian, A. V. Efremov, K. Goeke, A. Metz, P. Schweitzer and T. Teckentrup, Phys. Rev. D **77**, 014023 (2008).
47. P. Schweitzer *et al.*, Phys. Rev. D **64** (2001) 034013.
K. Goeke *et al.*, Acta Phys. Polon. B **32**, 1201 (2001).
M. Wakamatsu and T. Kubota, Phys. Rev. D **60**, 034020 (1999).
48. A. V. Efremov, O. V. Teryaev and P. Zavada, Phys. Rev. D **70** (2004) 054018.
P. Zavada, Eur. Phys. J. C **52**, 121 (2007), Phys. Rev. D **55**, 4290 (1997),
Phys. Rev. D **67**, 014019 (2003).
49. H. Avakian *et al.*, JLab E05-113, "Semi-Inclusive Pion Production with a Longitudinally Polarized Target at 6 GeV".
50. H. Avakian *et al.*, JLab PR12-07-107, "Studies of Spin-Orbit Correlations with Longitudinally Polarized Target".